

Cannabis, Cannabinoids, and Sleep: a Review of the Literature

Kimberly A. Babson¹ · James Sottile² · Danielle Morabito¹

Published online: 27 March 2017

© Springer Science+Business Media New York (outside the USA) 2017

Abstract

Purpose of Review The current review aims to summarize the state of research on cannabis and sleep up to 2014 and to review in detail the literature on cannabis and specific sleep disorders from 2014 to the time of publication.

Recent Findings Preliminary research into cannabis and insomnia suggests that cannabidiol (CBD) may have therapeutic potential for the treatment of insomnia. Delta-9 tetrahydrocannabinol (THC) may decrease sleep latency but could impair sleep quality long-term. Novel studies investigating cannabinoids and obstructive sleep apnea suggest that synthetic cannabinoids such as nabilone and dronabinol may have short-term benefit for sleep apnea due to their modulatory effects on serotonin-mediated apneas. CBD may hold promise for REM sleep behavior disorder and excessive daytime sleepiness, while nabilone may reduce nightmares associated with PTSD and may improve sleep among patients with chronic pain.

Summary Research on cannabis and sleep is in its infancy and has yielded mixed results. Additional controlled and longitudinal research is critical to advance our understanding of research and clinical implications.

Keywords Cannabis · Cannabinoids · Sleep · Insomnia · Sleep apnea

This article is part of the Topical Collection on *Sleep Disorders*

✉ Kimberly A. Babson
Kimberly.Babson@va.gov

¹ National Center for PTSD-Dissemination & Training Division, VA Palo Alto Health Care System, 795 Willow Road, Menlo Park, CA 94025, USA

² Palo Alto University, Palo Alto, CA, USA

Introduction

Rates of cannabis use within the USA continue to increase, with 8.3% of the US population reporting cannabis use within the past month [1]. Cannabis use has been associated with the development of cannabis use disorders, particularly among “at-risk” populations. In contrast, there has also been evidence to suggest that cannabis may have therapeutic potential. Indeed, as of January 1, 2017, 26 states and the District of Columbia have legalized cannabis for medical purposes, while 7 states and the District of Columbia have legalized the recreational use of cannabis. Understanding the research on both sides of this coin is important for clinical, research, and policy purposes.

The current paper seeks to provide a state-of-the-science review of the research on cannabis and sleep, a condition for which individuals often report using cannabis [2]. Previous review papers have provided an overview of the research on cannabis and sleep up through 2014. Therefore, within this review, we will provide a summary of the hallmark work in this area through 2014 along with an update of new research from 2014 to the end of 2016. First, we will provide a primer on cannabis and cannabinoids and how they relate to sleep. We will then provide an overview of research on both sides of this topic, namely, the risk and potential benefits of cannabis on sleep and the impact of poor sleep on cannabis use. Finally, we will provide an overview of the research on cannabis and specific sleep disorders. We will conclude with an integrative summary and call for future directions.

Introduction to Cannabinoids

The cannabis flower is comprised of over 100 different cannabinoids, the active compounds found within the cannabis plant. Cannabinoids work on the endocannabinoid system

(ECS) which consists of a series of neuromodulatory lipids and receptors located throughout the brain and central and peripheral nervous system, which accept endogenous cannabinoids (anandamide, 2-arachidonoylglycerol) and phytocannabinoids (plant-based). The CB1 and CB2 receptors are two main receptors within the ECS [3]. The two most well-researched phytocannabinoids are delta-9 tetrahydrocannabinol (THC) and cannabidiol (CBD). THC is the primary psychoactive component of cannabis which is responsible for the “high” associated with cannabis use. THC acts on CB1 receptors and yields a biphasic effect such that the impact of THC varies between low and high doses. CBD is a non-intoxicating constituent of cannabis which acts on CB2 receptors. CBD has been shown to counter the effects of THC and has received a lot of attention for its potential therapeutic effects [4].

Based on the potential therapeutic impact, cannabis-based medicine extracts have been developed. These extracts are synthetic THC (dronabinol, nabilone), CBD (Charlotte’s web), and nabiximols (1:1 CBD/THC, Sativex) which are delivered orally. The development of synthetic extracts has allowed for investigation of the effects of specific cannabinoids.

The Impact of Cannabis on Sleep

Research on the impact of cannabis on sleep started in the 1970s and included a number of studies examining polysomnography (PSG)-based sleep. This resulted in mixed findings with some work showing a decrease in sleep onset latency [5] and wake after sleep onset [6], while other work did not replicate these findings [7], but instead observed an increase in slow wave sleep [7, 8] and a decrease in REM [6, 7, 9]. Additional work from this era also suggested that cannabis may have a short-term benefit on sleep, particularly in reducing sleep onset latency [10]; however, chronic use of cannabis could be associated with habituation to the sleep inducing and slow wave sleep-enhancing properties [8, 11–13]. This initial work suggested that long-term use could have a negative impact on sleep in two primary ways. First, individuals may find themselves in a vicious cycle of using cannabis to manage sleep, habituating to the effects, and using more cannabis in order to obtain the desired impact, resulting in problematic patterns of use. Second, sleep disturbances are the hallmark of cannabis withdrawal and may serve to maintain use and predict relapse.

Sleep and Cannabis Relapse A breadth of evidence has converged to demonstrate that poor sleep is a critical risk factor for predicting cannabis cessation success [14–16]. Poor sleep quality prior to a quit attempt has been shown to increase the risk of early lapse/relapse to cannabis [16] and be associated with less of a reduction in cannabis use frequency among cannabis-dependent veterans [15]. Post-quit sleep has also

been shown to impact quit success and rates of lapse/relapse to use [14, 17–19]. For example, Budney and colleagues [14] demonstrated that 65% of cannabis users reported poor sleep as the primary reason for lapse/relapse to cannabis during a prior quit attempt.

Cannabis Withdrawal and Sleep Nonclinical and clinical research has now characterized the profile of cannabis withdrawal, with sleep disturbances and vivid dreams representing hallmark cannabis withdrawal symptoms [20]. Indeed disturbed sleep can last up to 45 days post-cessation making this the longest lasting withdrawal symptom [20]. Disturbed sleep is commonly reported with 67–73% of adults and 33–43% of adolescents reporting disturbed sleep during a quit attempt [14, 21, 22]. This work has combined to suggest that sleep disturbance is one of the most severe cannabis withdrawal symptoms [23]. While a majority of this work was initially based on self-reported sleep questionnaires, more recent work has demonstrated objective changes in sleep during cannabis withdrawal. In a cross-sectional study using PSG, Bolla and colleagues [24] demonstrated that abrupt cannabis cessation among heavy users was associated with a decrease in total sleep time, sleep efficiency, and %REM. In addition, increases in wake after sleep onset, sleep onset latency, and periodic limb movements were observed. Vandrey and colleagues [19] examined PSG-measured sleep in a sample of heavy cannabis users who completed a within-subject crossover study which alternated between periods of cannabis use and an abstinence phase supplemented by administration of either a placebo or zolpidem. Results demonstrated that abrupt cessation was associated with an increase in sleep onset latency and %REM, while a decline in sleep efficiency was observed. Administration of zolpidem attenuated the effects such that there was no difference in PSG sleep between the zolpidem and cannabis use phases. In addition, zolpidem reversed the abstinence-induced changes in stage 2 sleep and REM. Taken together, self-reported sleep and objective indices of poor sleep have been consistently demonstrated during cannabis withdrawal.

The Role of the Endocannabinoid System on the Circadian Sleep–Wake Cycle

The role of sleep in cannabis use and withdrawal is not surprising as recent work has demonstrated that the ECS is involved in the regulation of the circadian sleep–wake cycle [25], including the maintenance and promotion of sleep [26]. Specifically, it has been hypothesized that the ECS serves as the link between circadian regulation systems (i.e., superchiasmatic nucleus) and the behavioral and physiological processes that are affected, including sleep [26].

The role of the ECS on circadian rhythms has been further supported by work demonstrating that a lack of normal sleep

causes dysregulation within the ECS [26], while elevation in the ECS at the receptor level is involved in the homeostatic recovery of sleep after non-normal sleep [26]. Based on this backdrop, the ECS is a critical system involved in the regulation of the circadian rhythm sleep–wake cycle, highlighting the importance of examining the impact of cannabinoids on sleep.

Effects of Cannabinoids on the Sleep–Wake Cycle

The mid-2000s has seen resurgence in research focused on cannabis and sleep. In comparison to the work done in the 1970s, the majority of this work has focused on self-reported sleep and has also started to investigate the potential therapeutic impact of cannabinoids for sleep. This resurgence in research is likely due to an increased sophistication and understanding of cannabis and the constituent components. Indeed, recent research has demonstrated that the type of cannabinoids (THC, CBD), ratio of cannabinoids, dosage, timing of administration, and route of administration all play a critical role in outcomes.

The Role of THC on the Sleep–Wake Cycle Preclinical research has suggested that circadian rhythms are less pronounced (assessed by body temperature over 2 weeks) during THC administration [27]. In addition, within the context of early clinical research, chronic administration of THC has been shown to result in the development of tolerance to sleep effects [26]. Further work among adults ($n = 8$) has demonstrated that 15 mg of THC did not have an impact on nocturnal sleep and resulted in increased sleepiness and delayed sleep onset the following day as well as changes in mood and memory difficulty [28].

The Role of CBD on the Sleep–Wake Cycle Administration of CBD has been shown to have differential effects on sleep based on dose. Indeed, low-dose CBD has a stimulating effect, while high-dose CBD has a sedating effect. In a study among individuals with insomnia, results suggested that administration of 160 mg/day of CBD increased total sleep time and decreased the frequency of arousals during the night [29], while low-dose CBD has been associated with increased wakefulness [28, 30].

Summary

Taken together, research suggests that short-term use of cannabis may have a therapeutic impact on sleep, specifically related to sleep onset latency and slow wave sleep. However, long-term chronic use is associated with habituation to the sleep-enhancing benefits and is associated with increased risk for cannabis dependence. Sleep disruption (self-reported and objective) is a primary withdrawal symptom

from cannabis and may play a role in cannabis lapse/relapse during cessation attempts. While this work has been focused on the cannabis flower, recent work has suggested that specific cannabinoids (THC and CBD) have a differential impact on sleep. This is further influenced by the dosage, ratio of cannabinoids, timing of administration, and route of administration. Initial work examining specific cannabinoids suggests a potential therapeutic effect of high-dose CBD and low-dose THC for sleep. What follows is an overview of the impact of cannabis, and specific cannabinoids, on a number of sleep disorders for which this research has been examined (see Table 1 for an overview of studies).

Cannabinoids and Specific Sleep Disorders

Insomnia/Sleep Quality

Insomnia is defined as dissatisfaction with sleep quantity or quality associated with difficulty falling asleep, difficulty maintaining sleep throughout the night, and/or waking up early in the morning with an inability to return to sleep that causes significant distress or impairment in functioning [52]. General prevalence rates of insomnia have increased in recent years from 17.5% in 2002 to 19.2% of the adult US population, representing 46.2 million [53•].

Recent work in animal models has focused on the effect of CBD on sleep quality and the sleep–wake cycle [32, 33]. Chagas and colleagues found an increase in total percentage of sleep in rats after administration of mid-range and high-dose CBD injections as compared to placebo. The effects on REM varied by dosage such that high-dose CBD increased REM sleep latency on the day of administration and mid-range dose CBD decreased REM sleep latency the day after administration [33]. Meanwhile, Hsiao and colleagues found that CBD blocked anxiety-induced REM sleep suppression but had no effect on NREM sleep [32]. This work is further supported by a recent case report in which administration of CBD oil reduced insomnia symptoms and PTSD-related sleep disturbances [54•]. Together, these findings suggest that CBD may impact sleep quality through its anxiolytic effects.

However, recent work measuring sleep in young adults has suggested that CBD may decrease stage 3 sleep, when used in combination with THC [28]. On the other hand, THC and synthetic THC preparations have been associated with decreased sleep latency [28, 55]. However, in Gorelick and colleagues synthetic THC administration study, overall amount of nighttime sleep decreased over time suggesting a potential effect of tolerance [55].

Recent studies on whole plant cannabis and sleep quality are similarly conflicting. Studies of medicinal cannabis users have found that individuals commonly report using cannabis for insomnia [31, 35•]. Among medical cannabis users,

Table 1 Recent studies on cannabis and sleep

Study (primary author, year)	Sample	Cannabinoid or cannabis-based medicine	Control group	Dose(s)	Route/timing of administration	Sleep measure (self-report)	Sleep measure (PSG)	Results
Nicholson, 2004 [28]	Healthy young adults age (21–34) N = 8	THC CBD	Placebo	Insomnia 15 mg THC 10 mg THC/CBD 1:1 30 mg THC/CBD 1:1	Oral 10:00 PM	Stanford Sleepiness Scale; Sam-Perelli fatigue rating	EEG EOG EMG Sleep Latency Test	No effect of 15 mg THC on nocturnal sleep but increased sleepiness and decreased sleep latency the following morning Combination low and high doses resulted in decreased stage 3 sleep Combination high dose increase wakefulness Decreased sleep latency in both groups after cannabis use CBD blocked anxiety-induced REM sleep suppression but had no effect on NREM sleep Increase in total percentage of sleep in 10 and 40 mg/kg conditions Increase in REM sleep latency in light period on day of administration (40 mg/kg) Decrease REM sleep latency day after administration (10 mg/kg) Higher synthetic THC concentrations in the evening were associated with decreased sleep latency Overall amount of nighttime sleep decreased during study
Tringale, 2011 [31]	Medical cannabis users N = 147	Cannabis (flower)	No cannabis use; no sleep problems	Varied	Varied	Sleep Latency	None	Decreased sleep latency in both groups after cannabis use CBD blocked anxiety-induced REM sleep suppression but had no effect on NREM sleep Increase in total percentage of sleep in 10 and 40 mg/kg conditions
Hsiao, 2012 [32]	Male Wistar rats N = 28	CBD	Placebo	0.5 µg, 1.0 µg	CeA injection 20 min prior to light onset	None	BEG EMG	CBD blocked anxiety-induced REM sleep suppression but had no effect on NREM sleep Increase in total percentage of sleep in 10 and 40 mg/kg conditions
Chagas, 2013 [33]	Male Wistar rats N = 28	CBD	Placebo	2.5 mg/kg 10 mg/kg 40 mg/kg	Intraperitoneal injection 7:00 AM–8:00 AM	None	ECoG EMG	Increase in total percentage of sleep in 10 and 40 mg/kg conditions
Gorelick, 2013 [34]	Daily cannabis users N = 13	Marinol	None	20 mg single dose; 40–120 mg daily	Around-the-clock oral admin	MEQ; St. Mary's Hospital Sleep	None	Higher synthetic THC concentrations in the evening were associated with decreased sleep latency Overall amount of nighttime sleep decreased during study
Study (primary author, year) Belendiuk, 2015 [35]	Sample Medical cannabis users N = 163	Cannabinoid Cannabis (flower)	Control group None	Dose(s) Varied	Route/timing of administration Varied	Sleep measure (self-report) PSQI	Sleep measure (PSG) None	Results Participants reported using cannabis for insomnia (N = 81) and nightmares (N = 14)

Table 1 (continued)

Study (primary author, year)	Sample	Cannabinoid or cannabis-based medicine	Control group	Dose(s)	Route/timing of administration	Sleep measure (self-report)	Sleep measure (PSG)	Results
Ogeil, 2015 [36]	Alcohol and/or cannabis users N = 248	Cannabis (flower)	None	Varied	Varied	PSQI	None	Those using for nightmares preferred sativa strains Individuals with insomnia and greater sleep latency reported using higher CBD concentrations Sleep medication use was associated with use of lower THC concentration Poor sleep quality was associated with problematic alcohol and cannabis use Women had poorer sleep outcomes than men
Shannon, 2016 [37]	Female with PTSD, age 10	CBD	Patient before CBD	25 mg capsule 3 mg spray	Oral administration 6:00 PM capsule and 2-4× spray at varied times	Sleep Disturbance Scale for children	None	CBD oil reduced insomnia and sleep disturbances in the patient
Carley, 2002 [38]	Male Sprague-Dawley rats N = 11	THC Oleamide	Crossover design	Obstructive sleep apnea (OSA) THC = 0.1, 1.0, or 10 mg/kg Oleamide = 0.1, 1.0, 10 mg/kg Scatomin = 0.79 mg/kg Combination = 0.1 mg/kg THC = 0.79 mg/kg serotonin or 0.1 mg/kg oleamide	12 intraperitoneal injections 15 min prior to PSG			Reduction in apnea index among THC and oleamide conditions
Jumpertz, 2010 [39]	Patients with sleep apnea	OEA AEA 2-AG	Healthy control	N/A	N/A	None	Oronasal airflow; Respiratory distress index	Three times higher concentrations of OEA in sleep apnea group suggesting a role of endocannabinoids in regulating wakefulness associated with respiratory distress
Study (primary author, year) Prasad, 2013 [40]	Sample Adults with sleep apnea N = 17	Cannabinoid Dronabinol	Control group None	Dose(s) 2.5 mg titrated to 10 mg	Route/timing of administration Oral administration 30 min prior to bed	Sleep measure (self-report) None	Sleep measure (PSG) EEG EOG EMG	Results Reduction in apnea-hypopnea index from baseline to night 21

Table 1 (continued)

Study (primary author, year)	Sample	Cannabinoid or cannabis-based medicine	Control group	Dose(s)	Route/timing of administration	Sleep measure (self-report)	Sleep measure (PSG)	Results
Cailik, 2014 [41•]	Male Sprague-Dawley rats N = 24 Adults with sleep apnea N = 17	Dronabinol	Vehicle control	100 µg/5 µl 10 µg/5 µl	Injection in nodose ganglia	None	Oronasal airflow; pulse oximetry; body position EMG/g Piezoelectric strain gauge	Reduction in apneas and hypopneas in dronabinol condition Shift in EEG power toward delta and theta frequencies and strengthening of ultradian rhythms associated with increasing doses of dronabinol
Farabi, 2014 [42•]	Adults with sleep apnea N = 17	Dronabinol	None	2.5 mg titrated to 10 mg	Oral administration 30 min prior to bed	None	EEG	Reduction in reflex apneas in the dronabinol condition No Reduction in peripherally induced reflex apneas in the dronabinol condition
Cailik, 2014 [43•]	Male Sprague-Dawley rats N = 36	Dronabinol	Vehicle control	100 µg/5 µl	Injection in nodose ganglia	None	EMG/g Piezoelectric strain gauge	Reduction in REM sleep behavior disorder symptoms after CBD treatment
Cailik, 2016 [44•]	Male Sprague-Dawley rats N = 30	Dronabinol	Vehicle control	100 µg/5 µl 10 µg/3 µl 1 µg/3 µl 0.1 µg/3 µl	Injection in right lateral ventricle	None	EMG/g Piezoelectric strain gauge	Reduction in REM sleep behavior disorder symptoms after CBD treatment
Chagas, 2014 [45•]	Patients with RBD N = 4	CBD	None	REM sleep behavior disorder (RBD) 75 mg/day 300 mg/day	Oral	None	PSG	Reduction in REM sleep behavior disorder symptoms after CBD treatment
Study (primary author, year)	Sample	Cannabinoid	Control group	Dose(s)	Route/Route/timing of administration	Sleep measure (self-report)	Sleep measure (PSG)	Results
Fraser, 2009 [46]	Patients with treatment-resistant PTSD N = 47 Adult male inmates with serious mental illness N = 104	Nabilone	None	Nightmares 0.5 mg titrated up to a max of 4 mg daily	Oral administration 1 h prior to bed	Nightmare presence and intensity; hours of sleep	None	Reduction in nightmares after nabilone
Cameron, 2014 [47•]	Adult male inmates with serious mental illness N = 104	Nabilone	None	Mean initial dose 1.4 mg/day titrated to 4 mg/day	Oral	Sleep hours per night and nightmares per week	None	Reduction in nightmares and increase in hours of sleep per night after nabilone
Roitman, 2014 [48•]	Outpatients with PTSD N = 10	THC	None	2.5 mg 2×/day titrated to 5 mg 2×/day	Oral administration 1 h after waking up and 2 h prior to bed	PSQI NFO NES	None	Reduction in nightmares and improvement in sleep quality after THC
Jetty, 2015 [49•]	Male military personnel with PTSD N = 10	Nabilone	Placebo	0.5 mg titrated to 3 mg	Oral administration 1 h prior to bed	CAPS sleep items; PTSD dream rating scale; sleep diary	None	Reduction in nightmares in the nabilone condition
Ware, 2010 [50]	Fibromyalgia patients with insomnia N = 29	Nabilone	Amitriptyline	Sleep in pain conditions 0.5 mg 1 mg	Oral	ISI Leeds Sleep Evaluation	None	Greater improvement in sleep quality in the nabilone condition

Table 1 (continued)

Study (primary author, year)	Sample	Cannabinoid or cannabis-based medicine	Control group	Dose(s)	Route/timing of administration	Sleep measure (self-report)	Sleep measure (PSG)	Results
Study (primary author, year)	Sample	Cannabinoid	Control group	Dose(s)	Route/timing of administration	Sleep Measure (Self-report)	Sleep measure (PSG)	Results
Nicholson, 2004 [28]	Healthy young adults age (21–34) N = 8	THC CBD	Placebo	Excessive daytime sleepiness (EDS) 15 mg THC 10 mg THC/CBD 1:1 30 mg THC/CBD 1:1	Oral 10:00 PM	Stanford Sleepiness Scale Sam-Perelli fatigue rating	EEG EOG EMG Sleep latency	15 mg THC increased sleepiness and decreased sleep latency the following morning Combination high-dose increase wakefulness Patients who screened positive for THC evidenced significantly more excessive daytime sleepiness symptoms and were more likely to meet criteria for narcolepsy
Dzodzomenyo, 2015 [51]	Pediatric patients N = 383	THC	Negative THC screen	Varied	Varied	None	Multiple Sleep Latency Test	

THC delta-9 tetrahydrocannabinol, CBD cannabidiol, EEG electroencephalography, EOG electrooculography, EMG electromyography, Cea central nucleus of the amygdala, REM rapid eye movement sleep, NREM non-rapid eye movement sleep, ECoG electrocorticography, EMGgg genioglossus electromyogram, MEQ Morning-Eveningness Questionnaire, PSQ/Pittsburgh Sleep Quality Index, OEA oleoyl ethanolamide, AEA anandamide, 2-AG 2-arachidonoylglycerol, RBD REM behavior disorder, PTSD posttraumatic stress disorder, NFQ Nightmare Frequency Questionnaire-Revised, NES nightmare effects survey, CAPS Clinical Administered PTSD Scale, ISI Insomnia Severity Index

individuals with insomnia and greater sleep latency reported using cannabis with higher CBD concentrations. Additionally, in this sample, sleep medication use was associated with use of lower THC concentrations [35•]. Tringale and colleagues found that medicinal cannabis users, both with and without reported sleep problems, experienced decreased sleep latency after cannabis use [31]. However, in a recent study of self-identified alcohol and/or cannabis users, cannabis use was associated with poor sleep quality [36•]. Taken together, these mixed findings suggest that cannabinoid concentration, dose, and route of administration may have differential effects on sleep quality and insomnia symptoms.

Obstructive Sleep Apnea

Obstructive sleep apnea (OSA) is the most prevalent form of sleep-disordered breathing in the USA affecting 9% of American adults [56]. Standard treatment for OSA is a mechanical device that keeps the airway clear for unlabored breathing called a continuous positive airway pressure (CPAP) machine. While CPAP machines are an effective treatment for OSA, many patients are noncompliant due to the discomfort associated with treatment [57•]. A number of animal and human studies have been conducted to examine cannabinoids as potential therapeutic alternatives for treating OSA.

This program of research was initiated in animal models. In an initial study, Carley and colleagues [38] sought to determine the role of endogenous and exogenous cannabinoids in modulating respiration during sleep among Sprague-Dawley rats. Results showed that both the endocannabinoid oleamide and the exogenous cannabinoid THC reduced apneic events, providing initial evidence to suggest that the cannabinoid system may function to suppress the serotonin-mediated symptoms of OSA. Indeed, serotonin has been shown to have a tonic (excitatory) effect on the upper airway motoneurons which serve to maintain upper airway patency and reduce apnea [58]. Years later, Calik and colleagues [41•] investigated the efficacy of injections of dronabinol (synthetic THC) into the nodose ganglion in reducing serotonin-induced apneas among Sprague-Dawley rats. The authors found that dronabinol reduced serotonin-induced apneas and modulated upper airway muscles responsible for regulating breathing during sleep, providing preliminary data for the use of dronabinol in treating adults with OSA. A follow-up study was then conducted to investigate the mechanism of action of dronabinol in preventing induced reflex apneas among adult male Sprague-Dawley rats [43•]. The authors pretreated the rats with AM251 and AM630 and CB₁ and CB₂ receptor antagonists, respectively, to determine whether this prevented the efficacy of injecting dronabinol into the nodose ganglia to reduce apneas. Results suggested that dronabinol's effects in suppressing apneas are facilitated by its action at both CB₁ and

CB₂ receptors. Finally, in the most recent work, Calik and colleagues [44•] investigated the effect of intracerebroventricular injections of dronabinol in suppressing apneas among Sprague-Dawley rats. The authors found no significant reductions in apneas among the rats treated with intracerebroventricular injections of dronabinol, which suggests that the efficacy of dronabinol in suppressing apneas is facilitated by peripheral rather than central nervous system activity.

This basic research has now been extended into work among humans with OSA. In a preliminary study, Jumpertz and colleagues [39] studied the function of endocannabinoids including oleoylethanolamide (OEA), anandamide (AEA), and 2-arachidonyl-glycerol (2-AG) in patients with OSA compared to healthy controls. The authors found higher concentrations of OEA but not AEA or 2-AG among patients with OSA, which were associated with difficulty breathing. This finding suggests that endocannabinoids, specifically OEA, may function to protect the brain from the symptoms of sleep apnea. Additional research was then conducted to examine the impact of dronabinol among humans with OSA. Here, Prasad and colleagues [40] investigated the safety, tolerability, and efficacy of dronabinol in decreasing the severity of OSA symptoms among adults. The authors found that dronabinol was safe, well tolerated, and demonstrated efficacy in reducing apneas among adults with OSA in doses ranging from 2.5 to 10 mg daily. In a separate study, Farabi and colleagues [42•] examined the effects of dronabinol on objective measures of the sleep process in adults with OSA. The authors found that dronabinol was associated with a change in delta and theta frequencies and an increase in ultradian rhythms, which was correlated with improvement in apneas and a decrease in sleepiness.

Overall, initial research conducted in animal models and preliminary work in humans suggests that synthetic forms of THC may have a therapeutic potential in the treatment of sleep apnea in the short term. Long-term follow-up studies and controlled trials are needed.

REM Behavior Disorder

REM sleep behavior disorder (RBD) is a parasomnia in which individuals lose muscle rigidity during REM sleep accompanying nightmares and may act out behaviors linked with dreams [59]. Research on the impact of cannabinoids on RBD is relatively limited. However, to date, one study has examined this relation. Specifically, Chagas and colleagues [45•] investigated the efficacy of CBD, the non-intoxicating constituent of cannabis, in reducing symptoms of RBD among four adults with Parkinson's disease. The authors found that CBD suppressed behaviors associated with RBD and was tolerated well by all patients. However, controlled research among larger samples, as well as long-term prospective

studies, has not been conducted, limiting the conclusions that can be drawn.

Nightmares

Nightmares associated with posttraumatic stress disorder (PTSD) are often a residual symptom that remains difficult to treat despite improvements in other domains [60]. Prazosin, an alpha-adrenergic blocker, is the only current pharmacological treatment for PTSD-related nightmares [61]. However, military veterans are increasingly using cannabinoids to treat symptoms of PTSD including nightmares [62]. This has sparked some initial research to examine the impact of cannabinoids on nightmares. Here, Fraser [46] reported on a study examining the effect of nabilone, a synthetic form of THC, in managing nightmares linked with PTSD. The authors found that treatment with nabilone produced a reduction in nightmare presence and intensity and increased participants' hours of sleep per night. Additional research conducted with a short-term follow-up has suggested that nabilone has been well-tolerated and reduced nightmares in military service members [49] and male prison inmates [47]. Finally, an open-label pilot study investigating the use of THC in the treatment of PTSD, found that THC led to an improvement in sleep quality and a reduction in the frequency of nightmares. However, mild adverse effects were reported among some patients such as dry mouth, headache, and dizziness [48]. Additional, controlled trials among larger, more diverse samples with the inclusion of long-term follow-up are needed.

Sleep in Pain Conditions

Chronic pain is a substantial public health issue, which affects about 20% of adults and is expected to increase as advancements in medical science allow people to live longer [63]. Chronic pain impacts individuals' ability to get restful sleep. Recent work has started to examine the potential role of cannabinoids in addressing sleep disturbances in the context of chronic pain [63]. The majority of this work has examined the impact of Sativex, a 1:1 THC/CBD cannabis-based medicine extract, on sleep in chronic pain. A review of clinical trials demonstrated self-reported improvement in sleep quality among chronic pain patients treated with Sativex [64]. However, other studies have suggested that 1:1 THC/CBD preparations do not increase sleep duration despite producing subjective improvements in sleep quality [65]. Nabilone, a synthetic THC, has also been examined within this context. Specifically, Ware and colleagues [50] conducted a study investigating the effectiveness of nabilone, compared with amitriptyline, in improving sleep among patients with fibromyalgia. The authors found that patients in both conditions evidenced improvements in sleep; however, treatment with

nabilone was associated with greater improvements in sleep compared to amitriptyline.

Daytime Sleepiness

Excessive daytime sleepiness (EDS) is a common symptom characterized by the urge to fall asleep during the light hours of the day with a number of causes including certain medications, various medical conditions, psychiatric conditions, and sleep disorders such as narcolepsy and OSA. The negative consequences of EDS which include behavioral, attention, memory, and immune impairment can have a significant impact on quality of life [66].

Research on cannabinoids and EDS is limited. However, a couple of notable studies have emerged in the past few years. Dzodzomenyo and colleagues' retrospective study of drug screens found that patients who screened positive for THC, indicating recent cannabis use, presented with more EDS and were more likely to meet criteria for narcolepsy [51]. Conversely, experimental evidence indicates the potential role of CBD in managing somnolence [67]. In fact, recent work on sleep and early morning behavior demonstrates CBD's impact on wakefulness. In this study, CBD administered in a 1:1 ratio counteracted the sedative effect of THC [28].

Conclusions

The purpose of this paper was to provide a state of the science review of the basic and human research on the associations between cannabis and cannabinoids and sleep, including specific sleep disorders. Overall, a breadth of work has demonstrated that sleep problems increase risk for lapse/relapse to cannabis [15, 16], and disturbed sleep is a hallmark withdrawal symptom that can last months after a cessation attempt [20]. Research examining the impact of whole plant cannabis on sleep has yielded mixed findings, with some work showing that cannabis use is associated with a decrease in sleep onset latency [5] and wake after sleep onset [6], while other work has not yielded these effects but instead noted an increase in slow wave sleep [7, 8] and a decrease in REM [6, 7, 9]. These mixed results are likely due to the heterogeneous nature of whole plant cannabis.

Research in recent years has been marked by sophistication in understanding the component constituents of cannabis and how these cannabinoids may have a differential impact on sleep and sleep disorders. Here, research has combined to suggest that THC may have a short-term sleep benefit; however, chronic administration of THC is associated with habituation to the sleep-enhancing qualities [26, 55]. In addition, THC has been associated with a less pronounced circadian rhythm [27] and has been associated with daytime sleepiness, delayed sleep onset latency (particularly at high doses and

over time), and negative mood and memory alterations [28, 51]. However, preliminary research has suggested a potential therapeutic effect of dronabinol and nabilone (THC based medicine extracts) on OSA [38–40, 41, 42, 43, 44] and nightmares [46, 47, 48, 49], respectively.

Work examining CBD, a non-intoxicating cannabinoid with a low addiction profile, has demonstrated differential effects based on dose. Specifically, low doses of CBD have been shown to be stimulating [28, 30] and has been investigated within the context of EDS [67]. In comparison, medium- and high-dose CBD is sedating [28, 30] and has been examined in the context of a number of sleep disorders including insomnia. Here, initial basic research has suggested that medium-/high-dose CBD is associated with an increase in the percentage of total sleep [32, 33]. This is supported by a pilot study in humans showing that high-dose CBD was associated with improved sleep [28]; however, when combined with THC may result in a decrease in slow wave sleep [28]. A preliminary study has also suggested that CBD may suppress REM behavior disorder in individuals with Parkinson's disease [45].

The majority of the studies that have examined the impact of cannabinoids on sleep have actually been done within the context of chronic pain conditions. A review of phase I and II clinical trials has suggested that a 1:1 THC/CBD ratio (Sativex) has been associated with sleep improvements among patients with chronic pain conditions [64]. Outside of this area of research, the work examining cannabinoids' association with sleep has been relatively limited which makes drawing conclusions difficult at this early stage of research.

Taken together, the research on cannabinoids and sleep is in its infancy and studies are limited by small sample sizes, short-term follow-up, and a lack of controls [c.f. 60]. However, the role of the endocannabinoid system on the circadian regulation system demonstrates the theoretical connection between cannabinoids and sleep. This highlights the critical need for continued research in this area to examine these questions in large samples, with adequate controls and long-term follow-up assessments in order to understand the long-term effects, a major question which still remains. Future research is needed to continue to examine the impact of specific cannabinoids on sleep, including objective measures of sleep. Within these studies, it is critical to examine the impact of cannabinoid ratios, dose, the timing of the dose, and route of administration as the current work combines to indicate these factors result in differential outcomes.

Compliance with Ethical Standards

Conflict of Interest James Sottile and Danielle Morabito declare that they have no conflict of interest.

Kimberly A. Babson has received personal fees from Insys Therapeutics.

Human and Animal Rights and Informed Consent This article does not contain any studies with human or animal subjects performed by any of the authors.

References

Papers of particular interest, published recently, have been highlighted as:

- Of importance
- Of major importance

1. Marijuana (Cannabis): Substance Abuse and Mental Health Services Administration; 2016 [Available from: <https://www.samhsa.gov/atod/marijuana>.
2. Bonn-Miller MO, Boden MT, Bucossi MM, Babson KA. Self-reported cannabis use characteristics, patterns and helpfulness among medical cannabis users. *Am J Drug Alcohol Abuse*. 2014;40(1):23–30. **This study examined medical cannabis patients' demographic profiles, reported reasons for using cannabis, and patterns of use.**
3. Pacher P, Batkai S, Kunos G. The endocannabinoid system as an emerging target of pharmacotherapy. *Pharmacol Rev*. 2006;58(3):389–462.
4. Grotenhermen F, Russo E. Cannabis and cannabinoids: pharmacology, toxicology, and therapeutic potential. New York: Haworth Press Inc; 2002.
5. Cousens K, DiMascio A. (–) Delta 9 THC as a hypnotic. An experimental study of three dose levels. *Psychopharmacologia*. 1973;33(4):355–64.
6. Pivik RT, Zarcone V, Dement WC, Hollister LE. Delta-9-tetrahydrocannabinol and synhexl: effects on human sleep patterns. *Clin Pharmacol Ther*. 1972;13(3):426–35.
7. Feinberg I, Jones R, Walker JM, Cavness C, March J. Effects of high dosage delta-9-tetrahydrocannabinol on sleep patterns in man. *Clin Pharmacol Ther*. 1975;17(4):458–66.
8. Barratt ES, Beaver W, White R. The effects of marijuana on human sleep patterns. *Biol Psychiatry*. 1974;8(1):47–54.
9. Feinberg I, Jones R, Walker J, Cavness C, Floyd T. Effects of marijuana extract and tetrahydrocannabinol on electroencephalographic sleep patterns. *Clin Pharmacol Ther*. 1976;19(6):782–94.
10. Chait LD. Subjective and behavioral effects of marijuana the morning after smoking. *Psychopharmacology*. 1990;100(3):328–33.
11. Freeman FR. The effect of chronically administered delta-9-tetrahydrocannabinol upon the polygraphically monitored sleep of normal volunteers. *Drug Alcohol Depend*. 1982;10(4):345–53.
12. Karacan I, Fernandez-Salas A, Coggins WJ, Carter WE, Williams RL, Thornby JI, et al. Sleep electroencephalographic-electrooculographic characteristics of chronic marijuana users: part I. *Ann N Y Acad Sci*. 1976;282:348–74.
13. Pranikoff K, Karacan I, Larson EA, Williams RL, Thornby JI, Hirsch CJ. Effects of marijuana smoking on the sleep EEG. Preliminary studies JFMA. 1973;60(3):28–31.
14. Budney AJ, Vandrey RG, Hughes JR, Thostenson JD, Bursac Z. Comparison of cannabis and tobacco withdrawal: severity and contribution to relapse. *J Subst Abus Treat*. 2008;35(4):362–8.
15. Babson KA, Boden MT, Bonn-Miller MO. The impact of perceived sleep quality and sleep efficiency/duration on cannabis use during a self-guided quit attempt. *Addict Behav*. 2013;38(11):2707–13.
16. Babson KA, Boden MT, Harris AH, Stickle TR, Bonn-Miller MO. Poor sleep quality as a risk factor for lapse following a cannabis quit attempt. *J Subst Abus Treat*. 2013;44(4):438–43.

17. Copersino ML, Boyd SJ, Tashkin DP, Huestis MA, Heishman SJ, Dermand JC, et al. Cannabis withdrawal among non-treatment-seeking adult cannabis users. *Am J Addict*. 2006;15(1):8–14.
18. Haney M. Self-administration of cocaine, cannabis and heroin in the human laboratory: benefits and pitfalls. *Addict Biol*. 2009;14(1):9–21.
19. Vandrey R, Smith MT, McCann UD, Budney AJ, Curran EM. Sleep disturbance and the effects of extended-release zolpidem during cannabis withdrawal. *Drug Alcohol Depend*. 2011;117(1):38–44.
20. Budney AJ, Moore BA, Vandrey RG, Hughes JR. The time course and significance of cannabis withdrawal. *J Abnorm Psychol*. 2003;112(3):393–402.
21. Crowley TJ, Macdonald MJ, Whitmore EA, Mikulich SK. Cannabis dependence, withdrawal, and reinforcing effects among adolescents with conduct symptoms and substance use disorders. *Drug Alcohol Depend*. 1998;50(1):27–37.
22. Vandrey R, Budney AJ, Kamon JL, Stanger C. Cannabis withdrawal in adolescent treatment seekers. *Drug Alcohol Depend*. 2005;78(2):205–10.
23. Vandrey RG, Budney AJ, Hughes JR, Liguori A. A within-subject comparison of withdrawal symptoms during abstinence from cannabis, tobacco, and both substances. *Drug Alcohol Depend*. 2008;92(1–3):48–54.
24. Bolla KI, Lesage SR, Gamaldo CE, Neubauer DN, Wang NY, Funderburk FR, et al. Polysomnogram changes in marijuana users who report sleep disturbances during prior abstinence. *Sleep Med*. 2010;11(9):882–9.
25. Sanford AE. Cannabinoids and hamster circadian activity rhythms. *Brain Res*. 2008;1222:141–8.
26. Vaughn LK, Denning G, Stuhr KL, de Wit H, Hill MN, Hillard CJ. Endocannabinoid signalling: has it got rhythm? *Br J Pharmacol*. 2010;160(3):530–43.
27. Perron RR, Tyson RL, Sutherland GR. Delta9-tetrahydrocannabinol increases brain temperature and inverts circadian rhythms. *Neuroreport*. 2001;12(17):3791–4.
28. Nicholson AN, Turner C, Stone BM, Robson PJ. Effect of Delta-9-tetrahydrocannabinol and cannabidiol on nocturnal sleep and early-morning behavior in young adults. *J Clin Psychopharmacol*. 2004;24(3):305–13.
29. Carlini EA, Cunha JM. Hypnotic and antiepileptic effects of cannabidiol. *J Clin Pharmacol*. 1981;21(8–9 Suppl):417S–27S.
30. Zuardi AW. Cannabidiol: from an inactive cannabinoid to a drug with wide spectrum of action. *Rev Bras Psiquiatr*. 2008;30(3):271–80.
31. Tringale R, Jensen C. Cannabis and insomnia. *Depression*. 2011;4(12):0–68.
32. Hsiao YT, Yi PL, Li CL, Chang FC. Effect of cannabidiol on sleep disruption induced by the repeated combination tests consisting of open field and elevated plus-maze in rats. *Neuropharmacology*. 2012;62(1):373–84.
33. Chagas MH, Crippa JA, Zuardi AW, Hallak JE, Machado-de-Sousa JP, Hirotsu C, et al. Effects of acute systemic administration of cannabidiol on sleep-wake cycle in rats. *J Psychopharmacol*. 2013;27(3):312–6.
34. Gorelick DA, Goodwin RS, Schwilke E, Schroeder JR, Schwoppe DM, Kelly DL, et al. Around-the-clock oral THC effects on sleep in male chronic daily cannabis smokers. *Am J Addict*. 2013;22(5):510–4.
35. Belendiuk KA, Babson KA, Vandrey R, Bonn-Miller MO. Cannabis species and cannabinoid concentration preference among sleep-disturbed medicinal cannabis users. *Addict Behav*. 2015;50:178. **This article documented the preferred cannabis strain and cannabinoid concentration among a group of medical cannabis users reporting use of cannabis for insomnia and nightmares.**
36. Ogeil R, Phillips JG, Rajaratnam SM, Broadbear JH. Risky drug use and effects on sleep quality and daytime sleepiness. *Hum Psychopharmacol Clin Exp*. 2015;30(5):356–63. **This study investigated sleep disturbance among a sample of problematic alcohol and cannabis users and found impaired sleep quality to be a larger problem than excessive daytime sleepiness.**
37. Shannon S, Opila-Lehman J. Effectiveness of cannabidiol oil for pediatric anxiety and insomnia as part of posttraumatic stress disorder: a case report. *Perm J*. 2016;20(4):108–11.
38. Carley DW, Paviovic S, Janelidze M, Radulovacki M. Functional role for cannabinoids in respiratory stability during sleep. *Sleep*. 2002;25(4):391–8.
39. Jumpertz R, Wiesner T, Bluhner M, Engeli S, Batkai S, Wirtz H, et al. Circulating endocannabinoids and N-acyl-ethanolamides in patients with sleep apnea—specific role of oleoylethanolamide. *Exp Clin Endocrinol Diabetes*. 2010;118(9):591–5.
40. Prasad B, Radulovacki MG, Carley DW. Proof of concept trial of dronabinol in obstructive sleep apnea. *Front Psychiatry*. 2013;4:1.
41. Calik MW, Radulovacki M, Carley DW. Intranodose ganglion injections of dronabinol attenuate serotonin-induced apnea in Sprague-Dawley rat. *Respir Physiol Neurobiol*. 2014;190:20–4. **This study examined the effect of synthetic cannabinoid injections into the nodose ganglion of rats and found that this treatment reduced symptoms of OSA.**
42. Farabi SS, Prasad B, Quinn L, Carley DW. Impact of dronabinol on quantitative electroencephalogram (qEEG) measures of sleep in obstructive sleep apnea syndrome. *J Clin Sleep Med*. 2014;10(1):49–56. **This study investigated dronabinol's effect on delta and theta wave activity and ultradian rhythms as measured by polysomnography.**
43. Calik MW, Radulovacki M, Carley DW. A method of nodose ganglia injection in Sprague-Dawley rat. *J Vis Exp*. 2014(93):e52233. **This article describes a technique for localized injection into the nodose ganglia of rats to prevent central nervous system alteration and isolate peripheral nervous system activity.**
44. Calik MW, Carley DW. Intracerebroventricular injections of dronabinol, a cannabinoid receptor agonist, does not attenuate serotonin-induced apnea in Sprague-Dawley rats. *J Negat Results Biomed*. 2016;15:8. **This study examined the effect of central nervous system injections of a synthetic cannabinoid and determined that the drug's ability to reduce apneas is a result of peripheral rather than central nervous system activation.**
45. Chagas MH, Eckeli AL, Zuardi AW, Pena-Pereira MA, Sobreira-Neto MA, Sobreira ET, et al. Cannabidiol can improve complex sleep-related behaviours associated with rapid eye movement sleep behaviour disorder in Parkinson's disease patients: a case series. *J Clin Pharm Ther*. 2014;39(5):564–6. **This article documented cannabidiol's effect in reducing REM sleep behavior disorder symptoms among a sample of patients with Parkinson's disease.**
46. Fraser GA. The use of a synthetic cannabinoid in the management of treatment-resistant nightmares in posttraumatic stress disorder (PTSD). *CNS Neurosci Ther*. 2009;15(1):84–8.
47. Cameron C, Watson D, Robinson J. Use of a synthetic cannabinoid in a correctional population for posttraumatic stress disorder-related insomnia and nightmares, chronic pain, harm reduction, and other indications: a retrospective evaluation. *J Clin Psychopharmacol*. 2014;34(5):559–64. **This study examined the use of Nabilone in a correctional facility and found that the drug reduced nightmares associated with PTSD.**
48. Roitman P, Mechoulam R, Cooper-Kazaz R, Shalev A. Preliminary, open-label, pilot study of add-on oral Delta9-tetrahydrocannabinol in chronic post-traumatic stress disorder. *Clin Drug Investig*. 2014;34(8):587–91. **This article provided preliminary evidence suggesting that THC may be effective in reducing nightmares and improving sleep quality among patients with PTSD.**
49. Jetly R, Heber A, Fraser G, Boisvert D. The efficacy of nabilone, a synthetic cannabinoid, in the treatment of PTSD-associated nightmares: a preliminary randomized, double-blind, placebo-controlled

- cross-over design study. *Psychoneuroendocrinology*. 2015;51:585–8. **This article provided preliminary evidence suggesting that Nabilone may be effective in reducing nightmares associated with PTSD.**
50. Ware MA, Fitzcharles MA, Joseph L, Shir Y. The effects of nabilone on sleep in fibromyalgia: results of a randomized controlled trial. *Anesth Analg*. 2010;110(2):604–10.
 51. Dzodzomenyo S, Stolfi A, Splaingard D, Earley E, Onadeko O, Splaingard M. Urine toxicology screen in multiple sleep latency test: the correlation of positive tetrahydrocannabinol, drug negative patients, and narcolepsy. *J Clin Sleep Med*. 2015;11(2):93–9. **This study investigated the relationship between positive urine screen for THC and presence of excessive daytime sleepiness and narcolepsy among a pediatric patient population.**
 52. Association AP. *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington 2013.
 53. Ford ES, Cunningham TJ, Giles WH, Croft JB. Trends in insomnia and excessive daytime sleepiness among U.S. adults from 2002 to 2012. *Sleep Med*. 2015;16(3):372–8. **This study examined prevalence of insomnia and daytime sleepiness based on National Health Interview survey data.**
 54. Shannon S, Opila-Lehman J. Cannabidiol oil for decreasing addictive use of marijuana: a case report. *Integr Med (Encinitas)*. 2015;14(6):31–5. **This case study investigated cannabidiol's effect on problematic cannabis use, anxiety, and sleep disturbance and found benefits for all three conditions.**
 55. Gorelick DA, Goodwin RS, Schwilke E, Schwoppe DM, Darwin WD, Kelly DL, et al. Tolerance to effects of high-dose oral delta9-tetrahydrocannabinol and plasma cannabinoid concentrations in male daily cannabis smokers. *J Anal Toxicol*. 2013;37(1):11–6.
 56. Peppard PE, Young T, Barnett JH, Palta M, Hagen EW, Hla KM. Increased prevalence of sleep-disordered breathing in adults. *Am J Epidemiol*. 2013;177(9):1006–14.
 57. Weaver TE, Calik MW, Farabi SS, Fink AM, Galang-Boquiren MT, Kapella MC, et al. Innovative treatments for adults with obstructive sleep apnea. *Nat Sci Sleep*. 2014;6:137–47. **This article provided a review of novel treatment approaches for OSA including surgical, mechanical, and pharmacotherapy treatments.**
 58. Veasey SC, Panckeri KA, Hoffman EA, Pack AI, Hendricks JC. The effects of serotonin antagonists in an animal model of sleep-disordered breathing. *Am J Respir Crit Care Med*. 1996;153(2):776–86.
 59. McCarter SJ, Boswell CL, St Louis EK, Dueffert LG, Slocumb N, Boeve BF, et al. Treatment outcomes in REM sleep behavior disorder. *Sleep Med*. 2013;14(3):237–42.
 60. Aurora RN, Zak RS, Auerbach SH, Casey KR, Chowdhuri S, Karippot A, et al. Best practice guide for the treatment of nightmare disorder in adults. *J Clin Sleep Med*. 2010;6(4):389–401.
 61. Raskind MA, Peterson K, Williams T, Hoff DJ, Hart K, Holmes H, et al. A trial of prazosin for combat trauma PTSD with nightmares in active-duty soldiers returned from Iraq and Afghanistan. *Am J Psychiatry*. 2013;170(9):1003–10.
 62. Betthausen K, Pilz J, Vollmer LE. Use and effects of cannabinoids in military veterans with posttraumatic stress disorder. *Am J Health Syst Pharm*. 2015;72(15):1279–84. **This study investigated the use of cannabinoids by military veterans with PTSD and found that cannabis use was associated with a decrease in nightmares associated with PTSD.**
 63. Lynch ME, Ware MA. Cannabinoids for the treatment of chronic non-cancer pain: an updated systematic review of randomized controlled trials. *J NeuroImmune Pharmacol*. 2015;10(2):293–301. **This article provided a literature review of research investigating the use of cannabinoids in treating chronic pain.**
 64. Russo EB, Guy GW, Robson PJ. Cannabis, pain, and sleep: lessons from therapeutic clinical trials of Sativex, a cannabis-based medicine. *Chem Biodivers*. 2007;4(8):1729–43.
 65. Ferguson G, Ware M. Review article: sleep, pain and cannabis. *Journal of Sleep Disorders & Therapy*. 2015;4(2):191. **This paper reviewed the state of current research on cannabis, chronic pain, and sleep.**
 66. Roth T. Effects of excessive daytime sleepiness and fatigue on overall health and cognitive function. *J Clin Psychiatry*. 2015;76(9):e1145. **This article described the negative health effects of excessive daytime sleepiness and discussed psychometric scales used to measure excessive daytime sleepiness.**
 67. Murillo-Rodriguez E, Sarro-Ramirez A, Sanchez D, Mijangos-Moreno S, Tejada-Padron A, Poot-Ake A, et al. Potential effects of cannabidiol as a wake-promoting agent. *Curr Neuropharmacol*. 2014;12(3):269–72. **This study examined the effects of cannabidiol as a wakefulness promoting drug.**